

CLENPIQ Instructions for Colonoscopy

Patient: _____ Date: _____ Admit time: _____

Preparing For Your Exam

Five days prior to your procedure:

- **STOP** your medications that have blood thinning effects. Examples: **Motrin, Ibuprofen, Aleve, Celebrex, and arthritis medications.** Tylenol and Aspirin are okay. (If you are on Coumadin, Plavix, Xarelto or any other anticoagulant/antiplatelet medication you must contact your Primary Care Physician for instructions.)
- **Diabetics:** If you are diabetic and take insulin, call your Primary Care Physician for instructions. Please check your blood sugar/glucose before leaving your home.

One day before your procedure:

- **NO FOOD all day!** Drink only CLEAR LIQUIDS (see below) starting when you wake up in the morning, throughout the entire day and up until 2 hours before your scheduled admit time. The more clear liquids you drink, the better the prep solution will work and the better you will feel. Below is a list of clear liquids that you may have. Please note that no red dye is allowed (no red Jell-O, red drinks, etc.). Please limit all dark beverages such as tea, soda, coffee, etc. to 2 cups daily.

Examples of Clear Liquids (NO RED DYE)

<u>Soda</u>	<u>Juice</u>	<u>Sports drinks</u>	<u>Other Options</u>	
All Sodas	Apple	Gatorade	Coffee/Tea (limit 2)	Water
Root Beer (limit 2)	Cranberry	Powerade	Crystal Light	Popsicle
Sprite/7UP	Grape	All Sport	Jell-O	
Coke/Pepsi (limit 2)	White Grape	Propel	Broth/Bouillon	

Day before your procedure at 5:00 PM:

- Drink one 5.4 oz bottle of CLENPIQ. Drink right from the bottle. It does not need to be mixed or diluted. Do not refrigerate or freeze CLENPIQ.
- After prep is completed, drink a minimal of 40oz of clear liquids over the next 5 hours.
- **Drink plenty of clear liquids thereafter.** Again, the more you drink, the better the prep works and the better you will feel.

***** If your prep is not working by 9:00 PM or if you are vomiting, please call physician at 402-441-5600*****

Day of your procedure:

- Take prep again at 5:00 AM if you have a morning procedure. Take prep at 8:00 AM if you have an afternoon procedure. Even if your prep results are already clear/yellow, you **WILL** still need to take the morning prep!!
- Drink the other 5.4oz bottle of CLENPIQ, followed by a minimal of 24 oz of clear liquids. **Drink clear liquids until two hours prior to your scheduled admit time**, then nothing more to drink.
- After completion of your second prep, your stools should look clear/yellow (urine like) and you should be able to see through the liquid to the bottom of the toilet.
- You may:
 1. Take your morning heart, blood pressure or seizure medication with sips of water.
 2. Brush your teeth and gargle in the morning.
 3. Do **NOT** smoke, chew tobacco, chew gum or suck on hard candy!

Please show up for your procedure at the admit time written at the top of this sheet.

*****Due to the sedation, you MUST have a responsible person with you and they MUST stay in our facility or your test will be cancelled. Taxi or bus rides are not allowed unless accompanied by a responsible adult.*****